

**Haku®**

EMULSIFIABLE LIQUID

TECHNICAL DATASHEET

FICHA TÉCNICA

## PERCENT COMPOSITION

% IN WEIGHT

## ACTIVE INGREDIENT

Oils of vegetable and animal origin

**Oil of oregano (*Lippia graveolens*)**  
(equivalent to 4.474 g a.i./L at 20°C)

0.50

**Peppermint oil (*Mentha piperita*)**

(equivalent to 8.948 g a.i./L at 20°C)

1.00

**Cinnamon oil (*Cinnamomum zeylanicum*)**

(equivalent to 13.422 g a.i./L at 20°C)

1.50

**Clove oil (*Syzygium aromaticum*)**

(equivalent to 8.948 g a.i./L at 20°C)

1.00

**Soybean oil (*Glycine max*)**

(equivalent to 40.266 g a.i./L at 20°C)

4.50

**Neem oil (*Azadirachta indica*)**

(equivalent to 58.162 g a.i./L at 20°C)

6.50

**Fish oil (mackerel)**

(equivalent to 134.22 g a.i./L at 20°C)

15.00

## INERT INGREDIENTS

Conditioners, Emulsifiers, Stabilizers and Diluents

70.00

TOTAL

100.00

RSCO-MEZC-INAC-1101Q-0680-096-30.0

OMRI. anm-18746

STOP, READ LABEL BEFORE USING PRODUCT MAY BE  
HARMFUL IF IN CONTACT WITH SKIN.

## METHODS FOR PREPARING AND APPLYING THE PRODUCT.

Shake the product well. Carefully open the container, measure the indicated amount of Haku® with a plastic measuring cylinder or any other specific graduated utensil for this use and pour it into a container containing half of its capacity of clean water. Shake and then add it to the application tank or spray backpack and shake again.

Haku® is an emulsifiable liquid, useful for the control of the pest listed below:

CROP	PEST	DOSE (L/ha)	NOTES
Blackberry Strawberry Blueberry (0)	Red spider mite ( <i>Tetranychus urticae</i> )	1.25 – 1.50	Two foliar applications at a 5-day interval Application volume 200-300 L water/ha.

(0): **Safety interval:** days from the last application to harvest.

**TIME OF RE-ENTRY INTO TREATED AREAS:** Once the spray has dried.

Av. Periférico Sur 3325 Piso 2, Co., San Jerónimo Lídice,  
Del. Magdalena Contreras C.P. 10200, CDMX

[www.vidra.bio](http://www.vidra.bio)