



Dilong® | MULTISIFIABLE LIQUID

Technical Data Sheet

PERCENTAGE COMPOSITION

ACTIVE INGREDIENT:

% BY WEIGHT

Polyphenols	0.400
Cinnamon Oil (<i>Cinnamomum zeylanicum</i>) (equivalent to 4.418 g a.i./L at 20°C)	
Black Pepper Oil (<i>Piper nigrum</i>) (equivalent to 4.418 g a.i./L at 20°C)	0.400
Neem Oil (<i>Azadirachta indica</i>) (equivalent to 13.254 g a.i./L at 20°C)	1.200
Polyphenols	1.500
Chaparral root extract (<i>Larrea tridentata</i>) (equivalent to 16.567 g a.i./L at 20°C)	
Cempasúchil flower extract (<i>Tagettes erecta</i>) (equivalent to 16.567 g a.i./L at 20°C)	1.500
Alkamides	0.125
Chaparral root extract (<i>Larrea tridentata</i>) (equivalent to 1.380 g a.i./L at 20°C)	
Ginger extract (<i>Zingiber officinale</i>) (equivalent to 0.276 g a.i./L at 20°C)	0.025
Ricinoleics	1.000
Higuerilla Oil (<i>Ricinus communis</i>) (equivalent to 11.045 g a.i./L at 20°C)	
Polycarboxylic Compounds	16.660
Chaparral root extract (<i>Larrea tridentata</i>) (equivalent to 184.00 g a.i./L at 20°C)	
Ginger extract (<i>Zingiber officinale</i>) (equivalent to 11.045 g a.i./L at 20°C)	3.340

Inert ingredients:

Conditioners, emulsifier, antifoaming agent, and diluent 73.850

TOTAL 100

RSCO-MEZC-INAC-01045-0551-096-26.15

**STOP! READ THE LABEL BEFORE USING THE PRODUCT
MAY BE HARMFUL IF SWALLOWED OR IN CONTACT WITH SKIN.**

DIRECTIONS FOR USE

Shake the product thoroughly before use. Carefully open the container, measure the required amount of Dilong® using a plastic graduated cylinder or measuring utensil, and pour it into a container filled halfway with clean water. Stir, then pour into the sprayer tank or knapsack sprayer and agitate again before application.

Av. Periférico Sur 3325 Piso 3, Col. San Jerónimo Lídice,
La Magdalena Contreras, C.P. 10200, CDMX

www.vidra.bio

During handling, preparation, and application, use appropriate personal protective equipment. Always calibrate your application equipment.

APPLICATION RECOMMENDATIONS:

CROP	PEST	RATE (L/HA)	REMARKS
Zucchini, Pumpkin, Chayote, Melon, Cucumber, Watermelon (No limit)	Flower thrips (<i>Frankliniella occidentalis</i>)	2.0	Apply 3 foliar sprays at 7-day intervals. Spray volume: 250–350 L of water per hectare.
Blueberry, Strawberry, Raspberry, Currant, Kiwi, Grape, Blackberry (0)	Thrips (<i>Frankliniella occidentalis</i>)	1.0-2.0	Apply 3 foliar sprays at 7-day intervals, when first individuals of the pest are detected. Spray volume: 600–650 L of water per hectare.
Avocado (0)	Thrips (<i>Frankliniella occidentalis</i>)	1.0-2.0	Apply 3 foliar sprays at 7-day intervals, when first individuals of the pest are detected. Spray volume: 410–460 L of water per hectare.