



PERCENTAGE COMPOSITION BY WEIGHT	% BY WEIGHT
ACTIVE INGREDIENT	
<i>Beauveria bassiana</i> Conidium Equivalent to 195.68 g a.i./L at 20°C, with a minimum content of 1.0×10 ⁷ CFU/mL	20.00
<i>Nomurea rileyi</i> Conidium Equivalent to 195.68 g a.i./L at 20°C, with a minimum content of 1.0×10 ⁷ CFU/mL	20.00
Neem Oil (<i>Azadirachta indica</i>) Equivalent to 39.136 g a.i./L at 20°C	4.00
<i>Bacillus thuringiensis</i> Spores (<i>kurstaki</i> and <i>sraelensis</i> varieties) Equivalent to 97.84 g a.i./L at 20°C, with a minimum content of 1.0×10 ⁷ CFU/mL	10.00
INERT INGREDIENTS Conditioner, emulsifier and diluent	46.00
TOTAL	100.00

RSCO-INAC-0195-0636-096-54

**STOP! READ THE LABEL BEFORE USING THE PRODUCT
MAY BE HARMFUL IF SWALLOWED OR IN CONTACT WITH SKIN.**

ALWAYS CALIBRATE YOUR APPLICATION EQUIPMENT.

Shake the product thoroughly before use. Carefully open the container, measure the required amount of JINN® using a plastic graduated cylinder or measuring utensil, and pour it into a container filled halfway with clean water. Stir, then pour into the sprayer tank or knapsack sprayer and agitate again before application.

JINN® is an emulsifiable liquid useful for controlling the following pests:

CROP	PEST	RATE (L/HA)	REMARKS
Broccoli, Cabbage, Cauliflower, Brussels Sprouts (3)	Diamondback moth (<i>Plutella xylostella</i>)	1.5	Apply 3 foliar sprays at 7-day intervals. Spray volume: 300–400 L of water per hectare.
Coffee (3)	Coffee berry borer (<i>Hypothenemus hampei</i>)	0.75 – 1.25	Apply 4 foliar sprays at 7-day intervals. Spray volume: 350–450 L of water per hectare.
Strawberry, Blueberry, Raspberry, Currant, Blackberry (3)	White grub (<i>Phyllophaga sp.</i>)	1.0 – 2.0	Apply 3 drenches at crop establishment, directed at the base of the stem, at 15-day intervals. Recommended spray volume: 800 L of water per hectare.

(): Pre-harvest interval: Days between the last application and harvest.

Re-entry interval: Once the spray has dried.