



Agroshield® | AQUEOUS

FICHA TÉCNICA

COMPOSITION

Ingredient	% by weight	Additional information
Hydrogen tetraoxophosphate(V): phosphoric acid	10.00	Equivalent to 106.83 g a.i./L at 20°C
Inert ingredients: surfactant and diluent	90.00	
Total	100.00	

RSCO-INAC-0592-X0506-013-10.0

STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED, IN CONTACT WITH SKIN, OR IF INHALED.

PRODUCT DESCRIPTION

Agroshield® is an agricultural insecticide-miticide of miscellaneous origin.

PREPARATION AND APPLICATION

Always calibrate application equipment. Wear the personal protective equipment specified on the label. Prepare a premix in a partially filled spray tank, add the recommended rate, agitate thoroughly, add the remaining water, and apply the mixture the same day.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Avocado	Brown mite (<i>Oligonychus punicae</i>)	0.21-0.43 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 155.0-165.7 gal water/acre
Garlic, onion, green onion, asparagus, leek	Onion thrips; leafminer	0.16-0.27 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 37.4-48.1 gal water/acre
Eggplant, pepper, tomato, potato, bell pepper, tomatillo	Whitefly nymphs and adults	0.16-0.27 US gal/acre	0 days	3 foliar applications at 0, 5 and 7 days. 58.8-69.5 gal water/acre
Zucchini, squash, chayote, melon, cucumber, watermelon	Whitefly nymphs and adults	0.11-0.21 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 58.8-69.5 gal water/acre
Citron, lime, lemon, mandarin, orange, grapefruit, tangerine	Asian citrus psyllid nymphs and adults	0.16-0.27 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 58.8-69.5 gal water/acre
Papaya	Whitefly nymphs and adults	0.11-0.21 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 58.8-69.5 gal water/acre
Ornamental crops listed on source label	Greenhouse whitefly nymphs and adults	0.11-0.21 US gal/acre	0 days	3 foliar applications at 0, 4 and 7 days. 58.8-69.5 gal water/acre
Blueberry, raspberry, strawberry, currant, kiwi, grape, blackberry	Whitefly and two-spotted spider mite	0.16-0.27 US gal/acre	0 days	3 foliar applications at 4-day intervals. Source label states 4000-500 L/ha (427.6-534.5 gal water/acre).