



## Haku® | SOLUBLE CONCENTRATE

### FICHA TÉCNICA

#### COMPOSITION

| Ingredient                                              | % by weight | Additional information             |
|---------------------------------------------------------|-------------|------------------------------------|
| Eucalyptus extract (Eucalyptus globulus)                | 31.60       | Equivalent to 300 g a.i./L at 20°C |
| Coriander extract (Coriandrum sativum)                  | 10.53       | Equivalent to 100 g a.i./L at 20°C |
| Inert ingredients: solvents, stabilizers and surfactant | 57.87       |                                    |
| Total                                                   | 100.00      |                                    |

RSCO-MEZC-INAC-0930K-X0193-375-42.13

**STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED, IN CONTACT WITH SKIN, OR IF INHALED.**

#### PRODUCT DESCRIPTION

Haku® is a botanical insecticide-miticide formulated as a soluble concentrate.

#### PREPARATION AND APPLICATION

Wear complete personal protective equipment. Measure the required amount, prepare a premix in a partially filled spray tank, add the recommended rate, agitate thoroughly, add the remaining water, maintain agitation, and apply the mixture on the same day.

#### RECOMMENDED USES

| Crop(s)                                                  | Target pest/disease or use                      | Rate                  | Preharvest interval | Directions                                                         |
|----------------------------------------------------------|-------------------------------------------------|-----------------------|---------------------|--------------------------------------------------------------------|
| Avocado                                                  | Brown mite; avocado persea mite                 | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 58.8-69.5 gal water/acre |
| Blueberry, raspberry, strawberry, currant, blackberry    | Two-spotted spider mite                         | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 58.8-69.5 gal water/acre |
| Eggplant, pepper, tomato, bell pepper, tomatillo         | Two-spotted spider mite (protected agriculture) | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 42.8-53.5 gal water/acre |
| Eggplant, pepper, tomato, potato, bell pepper, tomatillo | Broad mite                                      | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 42.8-53.5 gal water/acre |
| Zucchini, squash, melon, cucumber, watermelon            | Two-spotted spider mite                         | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 42.8-53.5 gal water/acre |
| Asparagus                                                | Two-spotted spider mite                         | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 48.1-58.8 gal water/acre |
| Strawberry                                               | Cyclamen mite                                   | 0.11-0.21 US gal/acre | 0 days              | 3 foliar applications at 7-day intervals. 42.8-53.5 gal water/acre |
| Citrus crops                                             | Citrus red mite; citrus rust mite               | 0.11-0.21 US gal/acre | 0 days              | Apply at first detection following label directions.               |