



Jinn® | EMULSIFIABLE LIQUID

FICHA TÉCNICA

COMPOSITION

Ingredient	% by weight	Additional information
Beauveria bassiana conidia	20.00	At least 1.0×10^7 CFU/mL; equivalent to 195.68 g a.i./L
Nomuraea rileyi conidia	20.00	At least 1.0×10^7 CFU/mL; equivalent to 195.68 g a.i./L
Neem oil (Azadirachta indica)	4.00	Equivalent to 39.136 g a.i./L
Bacillus thuringiensis spores (kurstaki and israelensis varieties)	10.00	At least 1.0×10^7 CFU/mL; equivalent to 97.84 g a.i./L
Inert ingredients	46.00	Conditioner, emulsifier and diluent
Total	100.00	

RSCO-INAC-0195-0636-096-54

STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED OR IN CONTACT WITH SKIN.

PRODUCT DESCRIPTION

Jinn® is an emulsifiable liquid used to control the pests listed below.

PREPARATION AND APPLICATION

Always calibrate application equipment. Shake thoroughly, measure the required amount, add it to a container half-filled with clean water, agitate, then add to the spray tank or backpack sprayer and agitate again.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Broccoli, cabbage, cauliflower, Brussels sprouts	Diamondback moth (<i>Plutella xylostella</i>)	0.16 US gal/acre	3 days	3 foliar applications at 7-day intervals. 32.1-42.8 gal water/acre
Coffee	Coffee berry borer (<i>Hypothenemus hampei</i>)	0.08-0.13 US gal/acre	3 days	4 foliar applications at 7-day intervals. 37.4-48.1 gal water/acre
Strawberry, blueberry, raspberry, currant, blackberry	White grub (<i>Phyllophaga</i> sp.)	0.11-0.21 US gal/acre	3 days	3 drench applications to the stem base at establishment, 15 days apart. 85.5 gal water/acre