



Kauzar® | SOLUBLE CONCENTRATE

FICHA TÉCNICA

COMPOSITION

Ingredient	% by weight	Additional information
Quassia amara extract	9.64	Equivalent to 100 g a.i./L at 20°C
Mexican prickly poppy extract (Argemone mexicana)	4.82	Equivalent to 50 g a.i./L at 20°C
Castor oil (Ricinus communis)	4.82	Equivalent to 50 g a.i./L at 20°C
Inert ingredients: stabilizer, fortifier, solvent and surfactants	80.72	
Total	100.00	

RSCO-MEZC-INAC-0930U-X0202-375-19.28 | OMRI anm-22694

STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED, IN CONTACT WITH SKIN, OR IF INHALED.

PRODUCT DESCRIPTION

Kauzar® is a botanical insecticide formulated as a soluble concentrate.

PREPARATION AND APPLICATION

Always calibrate application equipment. Wearing complete personal protective equipment, open the container carefully, measure the required amount, prepare a premix in a partially filled spray tank, add the recommended rate, agitate thoroughly, add the remaining water, and maintain agitation. Apply the mixture on the same day it is prepared.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Garlic, onion, green onion, asparagus	Leafminer (<i>Liriomyza sativae</i>)	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. Begin at first detection. 48.1-58.8 gal water/acre
Celery, spinach, lettuce	Leafminer (<i>Liriomyza sativae</i>)	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. Begin at first detection. 37.4-48.1 gal water/acre
Celery, spinach, lettuce	Green peach aphid; potato aphid	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. Begin at first detection. 26.7 gal water/acre
Blueberry, raspberry, strawberry, currant, kiwi, grape, blackberry	Black bean aphid; whitefly; green peach aphid; cotton aphid	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. Begin at first detection. 37.4-58.8 gal water/acre
Eggplant, pepper, tomato, okra, potato, bell pepper, tomatillo	Leafminer; potato psyllid; whitefly; solenopsis mealybug	0.11-0.32 US gal/acre	0 days	3 foliar applications at 5- or 7-day intervals depending on pest. 48.1-69.5 gal water/acre
Broccoli, cabbage, cauliflower, Brussels sprouts, kohlrabi, canola, mustard	Cabbage aphid (<i>Brevicoryne brassicae</i>)	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. 26.7-37.4 gal water/acre
Zucchini, squash, chayote, melon, cucumber, watermelon	Whitefly; green peach aphid; leafminer	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. 37.4-53.5 gal water/acre
Papaya	Whitefly (<i>Bemisia tabaci</i>)	0.11-0.32 US gal/acre	0 days	3 foliar applications at 7-day intervals. 80.2-90.9 gal water/acre



Kauzar® | SOLUBLE CONCENTRATE

FICHA TÉCNICA

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Grape	Vine mealybug (Planococcus ficus)	0.11-0.32 US gal/acre	0 days	3 directed foliar applications at 7-day intervals. 48.1-58.8 gal water/acre