



Yasonte® | SUSPENSION CONCENTRATE

FICHA TÉCNICA

COMPOSITION

Ingredient	% by weight	Additional information
Bacillus amyloliquefaciens	30.30	1 × 10 ⁷ CFU/mL; equivalent to 300 g a.i./L at 20°C
Inert ingredients: surfactants, stabilizer, antifoam and solvent	69.70	
Total	100.00	

RSCO-FUNG-0395BA-X0717-064-30.30 | OMRI anm-22573

STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED, IN CONTACT WITH SKIN, OR IF INHALED.

PRODUCT DESCRIPTION

Yasonte® is a microbial-origin fungicide formulated as a suspension concentrate.

PREPARATION AND APPLICATION

Wear complete personal protective equipment. Measure the required amount, prepare a premix in a partially filled spray tank, add the recommended rate, agitate thoroughly, add the remaining water, maintain agitation, and apply the mixture on the same day.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Tomato, eggplant, pepper, bell pepper, tomatillo, potato	Powdery mildew; early blight	0.11-0.21 US gal/acre	0 days	3 foliar applications at 7-day intervals. 44.6-55.3 gal water/acre
Same crops	Dry rot (Fusarium oxysporum)	0.11-0.21 US gal/acre	0 days	3 drench applications directed to the plant crown at 7-day intervals. 50 mL mixture/plant; 171.0-181.7 gal water/acre
Same crops	Gray mold (Botrytis cinerea)	0.11-0.21 US gal/acre	0 days	3 foliar applications at 7-day intervals. 42.8-53.5 gal water/acre
Pepper, eggplant, tomato, okra, potato, bell pepper, tomatillo	Late blight / plant wilt (Phytophthora infestans)	0.11-0.21 US gal/acre	0 days	3 foliar applications at 7-day intervals, beginning 7-15 days after transplanting. 37.4 gal water/acre
Same crops	Damping-off (Fusarium, Verticillium, Rhizoctonia)	0.11-0.21 US gal/acre	0 days	3 drench applications to the plant crown at 7-day intervals after transplanting. 144.3-155.0 gal water/acre
Same crops	Vascular wilt (Fusarium oxysporum)	0.11-0.21 US gal/acre	0 days	3 drench applications to the plant crown at 7-day intervals. 176.4-187.1 gal water/acre
Zucchini, squash, chayote, melon, cucumber, watermelon	Damping-off; vascular wilt; powdery mildew; downy mildew	0.11-0.21 US gal/acre	0 days	3 drench or foliar applications at 7-day intervals according to disease. 37.4-155.0 gal water/acre
Blueberry, raspberry, strawberry, currant, blackberry	Powdery mildew; gray mold	0.11-0.21 US gal/acre	0 days	3 foliar applications at 7-day intervals. 48.1-58.8 gal water/acre