



TECHNICAL DATA SHEETS

Ardyan® | SOLUBLE CONCENTRATE

PERCENTAGE COMPOSITION

Ingredient	% by weight	Additional information
Aloe vera extract (<i>Aloe vera</i>)	68.46	Equivalent to 675 g a.i./L at 20°C
Pineapple extract (<i>Ananas comosus</i>)	22.82	Equivalent to 225 g a.i./L at 20°C
Antifoaming agent, solvent, and surfactant.	8.72	
Total:	100.00	

RSCO-MEZC-FUNG-0930R-X0191-375-91.28 | OMRI anm-22537

STOP, READ THE LABEL BEFORE USING THE PRODUCT.

MAY BE HARMFUL IF SWALLOWED. MAY BE HARMFUL IN CONTACT WITH SKIN. MAY BE HARMFUL IF INHALED.

PRODUCT DESCRIPTION

Ardyan® is a botanical fungicide formulated as a soluble concentrate.

PREPARATION AND APPLICATION

Always calibrate the application equipment. Wearing the adequate and complete protective equipment, open the container by unscrewing the cap until the exterior safety seal breaks and remove it completely. Invert the cap and place it on top of the container's opening, inserting the tab over the interior safety seal. Turn several times until this seal breaks. Perform this operation very carefully to avoid splashing and contact with skin and eyes. Measure the amount of product to be used with a plastic graduated cylinder or another specific graduated utensil for this purpose. Create a pre-mix with the necessary amount of product by filling the sprayer tank halfway with water, add the recommended dose of the product, agitate perfectly, and then add the remaining amount of water. Agitate continuously. Apply the mixture on the same day it is prepared.

RECOMMENDED USES

CROP(S)	DISEASE(S)	DOSAGE (L/ha)	SAFETY INTERVAL*	OBSERVATIONS
Blueberry Raspberry Strawberry Gooseberry Kiwi Grape Blackberry	Gray mold (<i>Botrytis cinerea</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 450 – 550 L of water/ha.
Raspberry	Yellow rust (<i>Pucciniastrum americanum</i> = <i>Aculeastrum americanum</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 550 – 650 L of water/ha.
Tomato Eggplant Chili pepper Potato Bell pepper Tomatillo	Gray mold (<i>Botrytis cinerea</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 450 – 550 L of water/ha.
Garlic Onion Welsh onion Leek	Gray mold (<i>Botrytis cinerea</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 450 – 550 L of water/ha.



TECHNICAL DATA SHEETS

Asparagus	Rust (<i>Puccinia asparagi</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 550 – 650 L of water/ha.
Lettuce	Gray mold (<i>Botrytis cinerea</i>)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 450 – 550 L of water/ha.
Barley	Stripe rust (<i>Puccinia striiformis</i> f. sp. hordei)	1.0 – 3.0	0	Perform 3 foliar applications at 7-day intervals. Begin at the first detection of disease symptoms and/or when environmental conditions are favorable for its development. Suggested application volume: 350 – 450 L of water/ha.

***Pre-Harvest Interval (PHI):** Days that must pass between the last application and harvest.

Re-entry interval for treated areas: Once the spray has dried.