



TECHNICAL DATA SHEETS

Haku® | SOLUBLE CONCENTRATE

PERCENTAGE COMPOSITION

Ingredient	% by weight	Additional information
Eucalyptus extract (<i>Eucalyptus globulus</i>)	31.60	Equivalent to 300 g a.i./L at 20°C
Cilantro extract (<i>Coriandrum sativum</i>)	10.53	Equivalent to 100 g a.i./L at 20°C
Inert ingredients: Stabilizers, surfactant, solvents.	57.87	
Total	100.00	

RSCO-MEZC-INAC-0930K-X0193-375-42.13 | OMRI anm-22524

STOP. READ THE LABEL BEFORE USING THIS PRODUCT.

MAY BE HARMFUL IF SWALLOWED. MAY BE HARMFUL IN CONTACT WITH SKIN. MAY BE HARMFUL IF INHALED.

PRODUCT DESCRIPTION

Haku® is a botanical insecticide formulated as a soluble concentrate.

PREPARATION AND APPLICATION

Always calibrate application equipment. Wearing full and appropriate personal protective equipment, open the container by unscrewing the cap until the outer safety seal breaks, then fully remove the cap. Invert the cap and place it on top of the container opening, inserting the tab into the inner safety seal; twist several times until the seal is broken. Perform this operation carefully to avoid splashing and contact with skin and eyes. Measure the required amount of product using a plastic graduated cylinder or any other specific measuring utensil. Prepare a pre-mix by filling half of the tank or sprayer reservoir with water, then add the recommended product dose. Mix thoroughly and add the remaining amount of water. Keep the mixture agitated at all times. Apply the solution on the same day it is prepared.

RECOMMENDED USES

CROP(S)	PEST(S)	DOSE (L/ha)	PRE-HARVEST INTERVAL (PHI)*	REMARKS
Avocado	Brown mite (<i>Oligonychus punicae</i>) Avocado bud mite (<i>Oligonychus perseae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first pest individuals. Suggested application volume: 550 – 650 L water/ha.
Blueberry Raspberry Strawberry Gooseberry Blackberry	Red spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, when the first pest individuals are detected. Suggested application volume: 550 – 650 L water/ha.
Eggplant Chili Tomato Bell pepper Green tomato	Red spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	In protected agriculture: Apply 3 foliar applications at 7-day intervals, when the first pest individuals are detected. Suggested application volume: 400 – 500 L water/ha.
Eggplant Chili Tomato Potato Bell pepper Green tomato	Broad mite (<i>Polyphagotarsonemus latus</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, when the first pest individuals are detected. Suggested application volume: 400 – 500 L water/ha.



TECHNICAL DATA SHEETS

Zucchini Squash Melon Cucumber Watermelon	Red spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first pest individuals. Suggested application volume: 400 – 500 L water/ha.
Asparagus	Red spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, when the first pest individuals are detected. Suggested application volume: 450 – 550 L water/ha.
Strawberry	Cyclamen mite (<i>Steneotarsonemus pallidus</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first pest individuals. Suggested application volume: 400 – 500 L water/ha.
Lemon Lime Mandarin Orange Grapefruit Pummelo	Citrus red mite (<i>Panonychus citri</i>) Citrus rust mite (<i>Phyllocoptruta oleivora</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, when the first pest individuals are detected. Suggested application volume: 400 – 500 L water/ha.
Apple Quince Pear Hawthorn	Two-spotted spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first infestations. Suggested application volume: 550 – 650 L water/ha.
Mango	Red spider mite (<i>Tetranychus urticae</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first pest individuals. Suggested application volume: 550 – 650 L water/ha.
Papaya	Red spider mite (<i>Tetranychus urticae</i>) Broad mite (<i>Polyphagotarsonemus latus</i>)	1.0 – 2.0	0	Apply 3 foliar applications at 7-day intervals, starting upon detection of the first pest individuals. Suggested application volume: 400 – 500 L water/ha.

***Pre-Harvest Interval (PHI):** Number of days that must pass between the last application and harvest.

RE-ENTRY INTERVAL FOR TREATED AREAS: After spray has completely dried.