



TECHNICAL DATA SHEETS

Jinn® | EMULSIFIABLE LIQUID

PERCENTAGE COMPOSITION

Ingredient	% by weight	Additional information
<i>Beauveria bassiana</i> conidia	20.00	At least 1.0×10^7 CFU/mL; equivalent to 195.68 g a.i./L
<i>Nomuraea rileyi</i> conidia	20.00	At least 1.0×10^7 CFU/mL; equivalent to 195.68 g a.i./L
Neem oil (<i>Azadirachta indica</i>)	4.00	Equivalent to 39.136 g a.i./L
<i>Bacillus thuringiensis</i> spores (<i>kurstaki</i> and <i>israelensis</i> varieties)	10.00	At least 1.0×10^7 CFU/mL; equivalent to 97.84 g a.i./L
Inert ingredients	46.00	Conditioner, emulsifier and diluent
Total	100.00	

RSCO-INAC-0195-0636-096-54

STOP. READ THE LABEL BEFORE USE. MAY BE HARMFUL IF SWALLOWED OR IN CONTACT WITH SKIN.

PRODUCT DESCRIPTION

Jinn® is an emulsifiable liquid used to control the pests listed below.

PREPARATION AND APPLICATION

Always calibrate application equipment. Shake thoroughly, measure the required amount, add it to a container half-filled with clean water, agitate, then add to the spray tank or backpack sprayer and agitate again.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Broccoli, cabbage, cauliflower, Brussels sprouts	Diamondback moth (<i>Plutella xylostella</i>)	1.5 L/ha	3 days	Make 3 foliar applications at 7-day intervals. Application volume: 300 – 400 L water/ha.
Coffee	Coffee berry borer (<i>Hypothenemus hampei</i>)	0.75 – 1.25 L/ha	3 days	Make 4 foliar applications at 7-day intervals. Application volume: 350 – 450 L water/ha.