



TECHNICAL DATA SHEETS

Dilong® | EMULSIFIABLE LIQUID

PERCENTAGE COMPOSITION

Ingredient	% by weight	Additional information
Cinnamon oil (<i>Cinnamomum zeylanicum</i>)	0.400	Equivalent to 4.418 g a.i./L
Black pepper oil (<i>Piper nigrum</i>)	0.400	Equivalent to 4.418 g a.i./L
Neem oil (<i>Azadirachta indica</i>)	1.200	Equivalent to 13.254 g a.i./L
Creosote bush root extract (<i>Larrea tridentata</i>)	1.500	Equivalent to 16.567 g a.i./L
Marigold flower extract (<i>Tagetes erecta</i>)	1.500	Equivalent to 16.567 g a.i./L
Creosote bush root extract (<i>Larrea tridentata</i>)	0.125	Equivalent to 1.380 g a.i./L
Ginger extract (<i>Zingiber officinale</i>)	0.025	Equivalent to 0.276 g a.i./L
Castor oil (<i>Ricinus communis</i>)	1.000	Equivalent to 11.045 g a.i./L
Creosote bush root extract (<i>Larrea tridentata</i>)	16.660	Equivalent to 184.00 g a.i./L
Ginger extract (<i>Zingiber officinale</i>)	3.340	Equivalent to 11.045 g a.i./L
Inert ingredients	73.850	Conditioners, emulsifier, antifoam and diluent
Total	100.000	

RSCO-MEZC-INAC-01045-0551-096-26.15 | OMRI anm-21809

STOP, READ THE LABEL BEFORE USING THIS PRODUCT.

MAY BE HARMFUL IF SWALLOWED. MAY BE HARMFUL IN CONTACT WITH SKIN. MAY BE HARMFUL IF INHALED.

PRODUCT DESCRIPTION

Dilong® is an emulsifiable liquid used to control the pests listed below.

PREPARATION AND APPLICATION

Shake thoroughly. Measure the required amount, add it to a container half-filled with clean water, agitate, then add to the spray tank or backpack sprayer and agitate again. Always calibrate application equipment and wear suitable personal protective equipment.

RECOMMENDED USES

Crop(s)	Target pest/disease or use	Rate	Preharvest interval	Directions
Zucchini, squash, chayote, melon, cucumber, watermelon	Western flower thrips (<i>Frankliniella occidentalis</i>)	2.0 L/ha	No limit	Make 3 foliar applications at 7-day intervals. Application volume: 250 – 350 L of water/ha.
Blueberry, strawberry, raspberry, currant, kiwi, grape, blackberry	Thrips (<i>Frankliniella occidentalis</i>)	1.0 – 2.0 L/ha	0 days	Make 3 foliar applications at 7-day intervals, starting at first detection. Application volume: 600 – 650 L of water/ha.
Avocado	Thrips (<i>Frankliniella occidentalis</i>)	1.0 – 2.0 L/ha	0 days	Make 3 foliar applications at 7-day intervals, starting at first detection. Application volume: 410 – 460 L of water/ha.